

CONTRIBUTION TO THE FLORA AND BIOGEOGRAPHY OF THE KIKLADES: FOLEGANDROS ISLAND (KIKLADES, GREECE)

K. KOUGIOUMOUTZIS, A. TINIAKOU, O. GEORGIU & T. GEORGIADIS

The island of Folegandros, located between the Milos and Santorini archipelagos in the southern Kiklades (Greece), constitutes together with Ios and Sikinos the south-central part of the phytogeographical region of the Kiklades. Its flora consists of 474 taxa, 47 of which are under statutory protection, 40 are Greek endemics and 145 are reported here for the first time. We show that Folegandros has the highest percentage of Greek endemics in the phytogeographical area of the Kiklades. The known distribution of the endemic *Muscari cycladicum* subsp. *cycladicum* is expanded, being reported for the first time outside the South Aegean Volcanic Arc. The floristic cross-correlation between Folegandros and other parts of the phytogeographical region of the Kiklades by means of Sørensen's index revealed that its phytogeographical affinities are stronger to Anafi Island than to any other part of the Kiklades.

Keywords. Aegean flora, biodiversity, endemism, phytogeography.

INTRODUCTION

The Aegean Sea has long attracted the attention of biogeographers (Turrill, 1929; Strid, 1996), since it is characterised by high environmental and topographical heterogeneity (Blondel *et al.*, 2010), diversity and endemism (Strid, 1996). The Aegean archipelago consists of more than 7000 islands and islets (Triantis & Mylonas, 2009), most of which are located in the phytogeographical region of the Kiklades. Intensive field work has taken place in this region, which is characterised as one of the most floristically explored phytogeographical regions of Greece (Dimopoulos *et al.*, 2013; Bergmeier & Strid, 2014). However, certain islands have been the main focal areas of floristic research, leaving our knowledge of the Kiklades' flora still incomplete.

Twenty-two large islands and numerous islets comprise the phytogeographical region of the Kiklades, which has long been considered floristically impoverished, compared to the other phytogeographical regions of Greece (Phitos *et al.*, 1995). The Kiklades host 1768 taxa (Dimopoulos *et al.*, 2013), 161 of which are Greek endemics (9.10%), and demonstrate lower than normal endemism, as expected by their size (Georghiou & Delipetrou, 2010). However, their degree of endemism is above average for Mediterranean islands (c.8%, Médail & Quézel, 1997) and in a recent analysis

(Kougioumoutzis & Tiniakou, 2014), some members of the Kiklades emerged as plant diversity hotspots. The Kiklades seem predestined to biogeographical and floristic studies, due to their variability in size, palaeogeographical history and geological composition, as well as in topographical and climatic characteristics. Less than half the large islands comprising the phytogeographical region of the Kiklades are floristically well known (Hansen, 1971; Tzanoudakis, 1981; Raus, 1986, 1988, 2012; Böhling, 1994; Browicz, 1997; Livaniou-Tiniakou *et al.*, 2003; Snogerup *et al.*, 2006; Kougioumoutzis *et al.*, 2012, 2014a). Thus, we decided to investigate the flora of Folegandros Island thoroughly.

The island of Folegandros (Fig. 1) has a special geographical, biogeographical and ecological position in the Aegean area. It is located in the southern part of the phytogeographical area of the Kiklades, between the volcanic archipelagos of Milos and Santorini, and, despite its small size (c.32 km²), it is characterised by a variety of substrates and is built up of five major tectono-stratigraphic units (Photiades & Keay, 2003), viz. schists with serpentinite bodies, gneiss, marbles, Quaternary and Neogene deposits. Folegandros forms a part of the median tectono-metamorphic Attic-Cycladic Complex, which is composed of an Alpine and pre-Alpine tectonic pile (Papanikolaou, 1984, 1987).

The study area is mainly hilly with an intense relief, the highest peak being Aghios Eleftherios hill (415 m). Several gravelly and sandy beaches can be found by the coast. The hydrographical network is rather limited, with no obvious runoff.




FIG. 1. Folegandros Island.

The nearest meteorological station to Folegandros lies in Antiparos island; according to Gouvas & Sakellariou (2011), this station and therefore the study area belongs to the arid bioclimatic zone with a mild winter and also to the Thermo-mediterranean zone, with a long dry period.

Rechinger (1944) was the first botanist to report on plants from Folegandros. Most records, however, were reported within the framework of the Natura 2000 research programme. Snogerup (1994), Runemark (1996, 2000, 2006), Strid & Tan (1997, 2002), Thanopoulos (2007) and Biel & Tan (2008) have also reported on plants from Folegandros. Information on some endemic taxa occurring in the area is given by Tan & Iatrou (2001). From a phytogeographical point of view, however, the interesting flora of the island of Folegandros has not yet received the attention it deserves.

The present study therefore aimed to thoroughly investigate the flora of Folegandros, by examining the floristic affinities of the study area to the other, floristically better explored islands of the phytogeographical region of the Kiklades, namely Anafi (Kougioumoutzis *et al.*, 2012), Andros (Snogerup *et al.*, 2006), Antiparos and Paros (Raus, 1986), Giaros (Tzanoudakis, 1981), Kimolos (Kougioumoutzis *et al.*, 2014a), Milos (Browicz, 1997; Raus, 2012), Naxos (Böhling, 1994), Santorini (Hansen, 1971; Raus, 1988; Tan & Iatrou, 2001) and Serifos (Livaniou-Tiniakou *et al.*, 2003).

MATERIALS AND METHODS

Several collection and field observation trips to the study area were carried out in the spring and autumn of 2013 in order to acquire an integrated knowledge of the flora and vegetation of Folegandros. Herbarium specimens are deposited at the Botanical Museum of the University of Patras (UPA). Species identification and nomenclature are according to Tutin *et al.* (1964–1980, 1993), Davis (1965–1985), Pignatti (1982), Tan & Iatrou (2001), Greuter *et al.* (1984–1989), Strid & Tan (1997, 2002), Greuter & Raab-Straube (2008) and Dimopoulos *et al.* (2013). Species identification and nomenclature of the genera *Anthemis* L., *Astragalus* L., *Anchusa* L., *Crepis* L., *Dittrichia* Greuter, *Reichardia* Roth, *Tordylium* L. and *Trifolium* L. are according to Georgiou (1990), Podlech (2008), Selvi & Bigazzi (2003), Kamari (1976), Brullo & de Marco (2000), Gallego *et al.* (1980), Al-Eisawi & Jury (1988) and Zohary & Heller (1984) respectively. For family delimitation we follow APG III (2009). The nomenclature and status of the endemic taxa recorded from Folegandros are based on Tan & Iatrou (2001), Georgiou & Delipetrou (2010) and Dimopoulos *et al.* (2013). The status of the alien taxa occurring in the study area is according to Arianoutsou *et al.* (2010) and Dimopoulos *et al.* (2013). The life-form and chorological categories follow Dimopoulos *et al.* (2013 – see Appendix for abbreviations used). Sørensen's index (Sørensen, 1948) as well as the statistical software SPSS 20 are used for the cross-correlation between the islands. We used the same databases as in Kougioumoutzis & Tiniakou (2014) and Kougioumoutzis *et al.* (2014b) regarding the endemism levels in the phytogeographical region of the Kiklades.

RESULTS

Flora

The vascular flora of Folegandros comprises 474 taxa, belonging to 263 genera and 68 families (Table 1). Six alien taxa are included in the plant list, but have not been considered in the floristic analysis.

The literature survey revealed 329 bibliographical reports for the study area (Rechinger, 1944; Snogerup, 1994; Runemark, 1996, 2000, 2006; Strid & Tan, 1997, 2002; Tan & Iatrou, 2001; Thanopoulos, 2007; Biel & Tan, 2008). We report 145 taxa as new to Folegandros (see Appendix). Forty taxa are Greek endemics, seven of which are new records for the study area. Fourteen of the new records and 47 taxa overall are under statutory protection.

The most species-rich families in the flora of Folegandros are the Asteraceae (64 taxa), followed by the Fabaceae (63 taxa) and Poaceae (48 taxa). These three families account for more than one third of the total flora (37.31%). Apiaceae (24 taxa), Caryophyllaceae (24 taxa), Brassicaceae (22 taxa) and Lamiaceae (14 taxa) are also well represented.

In life forms (Table 2) therophytes dominate, followed by hemicyptophytes, geophytes, chamaephytes and phanerophytes.

According to their general distribution, the local vascular flora can be classified into 15 main chorological groups (Table 3). The endemic group represents 8.55% of the total flora with 40 taxa. Phytogeographically, the endemic element is the most important group and is discussed separately. The Mediterranean chorological group predominates, highlighting the geographical position and climatic characteristics of the study area. The other chorological elements are represented in lower percentages, with a relatively high proportion of Asian elements, and also of invasive elements, indicating intense human impact in the study area.

The alien flora of Folegandros comprises six taxa (1.27%), belonging to six genera and six families. The neophytes amount to 60.00% of Folegandros alien flora and the most prominent among the invasive species is *Aeonium arboreum* (L.) Webb & Berthel., which occupies large areas, especially in the endemic species-rich calcareous cliffs of Folegandros.

TABLE 1. Number of vascular plant taxa in the flora of Folegandros Island

Systematic unit	Families	Genera	Taxa	%
Pteridophytes	3	5	5	1.06
Gymnospermae	2	2	2	0.42
Dicotyledones	49	200	374	78.90
Monocotyledones	14	56	93	19.62
Total	68	263	474	100.00

TABLE 2. Life forms in the flora of Folegandros Island

Life form	Total no. of taxa	%
Phanerophytes	24	5.13
Chamaephytes	37	7.91
Hemicryptophytes	86	18.38
Therophytes	269	57.48
Geophytes	51	10.90
Hydrophytes	1	0.20
Total	468	100.00

Endemism

According to Dimopoulos *et al.* (2013), 1768 taxa are found in the phytogeographical region of the Kiklades, 161 of which are considered endemics (9.10%). In Folegandros, 40 endemic taxa were found, making up 8.55% of its flora (Table 4). The level of endemism in Folegandros is rather low compared to that in the whole of Greece (22.2%, Dimopoulos *et al.*, 2013) but, taking into consideration the small size of the study area (c.32 km²), its geographic position not close to known areas of

TABLE 3. Chorological groups in the flora of Folegandros Island

Chorological group	No. of taxa	%	Total	
			No. of taxa	%
<i>1. Widely distributed taxa</i>			82	17.52
European-SW Asian	40	8.55		
Euro-Siberian	7	1.50		
Paleotemperate	15	3.20		
Circumtemperate	2	0.43		
Subtropical-Tropical	6	1.28		
Cosmopolitan	12	2.56		
<i>2. Mediterranean taxa</i>			339	72.43
E Mediterranean	53	11.32		
Mediterranean	197	42.09		
Mediterranean-Atlantic	8	1.71		
Mediterranean-European	35	7.48		
Mediterranean-SW Asian	46	9.83		
<i>3. Balkan taxa</i>			7	1.50
Balkan	1	0.21		
Balkan-Italian	2	0.43		
Balkan-Anatolian	4	0.86		
<i>4. Endemic taxa</i>			40	8.55
Endemic	40	8.55		
Total	468	100.00	468	100.00

TABLE 4. Endemism in Folegandros and the phytogeographical region of the Kiklades, as well as in the other large Kikladic islands

Region	% endemics
Kiklades	9.10
Amorgos	7.99
Anafi	5.99
Andros	4.04
Antiparos	3.85
Astypalaea	6.35
Folegandros	8.55
Giaros	2.52
Ios	3.67
Kea	3.36
Kimolos	6.88
Kythnos	5.44
Milos	5.54
Mykonos	3.48
Naxos	5.00
Paros	3.90
Polyaegos	3.70
Santorini	3.57
Serifos	4.26
Sifnos	5.48
Sikinos	6.71
Syros	2.44
Tinos	3.38

high endemism, the unfavourable arid climate, as well as the intense human pressure present on the island (i.e. high intensity grazing, highly cultivated areas – abandoned or not), this percentage is rather significant. Furthermore, compared to endemism levels in other parts of the phytogeographical region of the Kiklades, yet with larger size than that of the study area (Table 4), that of Folegandros is rather high, even appearing to be the highest in the phytogeographical region of the Kiklades.

The endemic species belong to 19 families and 31 genera. Families rich in endemic species in absolute numbers are Asteraceae, Campanulaceae and Caryophyllaceae (Table 5), their degree of endemism (9.38%, 80.00% and 16.67%, respectively) being higher than that of the general flora (8.55%). These results agree with the trend observed in the whole Greek endemic flora (Georghiou & Delipetrou, 2010).

Nearly half (19) of the endemic taxa found on Folegandros correspond to one or two phytogeographical areas (Table 6), thus providing valuable information regarding the phytogeographical position of the study area, as the existence of biregional endemics is a good indication of phytogeographical connections between regions (Georghiou & Delipetrou, 2010). It would be expected that Folegandros shows higher affinities with the phytogeographical area of the East Aegean Islands (EAe) since, according to Georghiou & Delipetrou (2010), the phytogeographical area of the

TABLE 5. Families with endemic taxa and their degree of endemism

Family	No. of endemic taxa	%
Asteraceae	6	9.38
Campanulaceae	4	80.00
Caryophyllaceae	4	16.67
Apiaceae	3	12.50
Brassicaceae	3	13.64
Iridaceae	3	60.00
Asparagaceae	2	25.00
Fabaceae	2	3.17
Plumbaginaceae	2	33.33
Ranunculaceae	2	25.00
Amaryllidaceae	1	10.00
Boraginaceae	1	7.14
Chenopodiaceae	1	10.00
Liliaceae	1	33.33
Orchidaceae	1	9.09
Poaceae	1	2.08
Primulaceae	1	25.00
Rubiaceae	1	7.69
Scrophulariaceae	1	10.00

Kyklades (Kik) is chorologically more closely connected to EAe than to that of Kriti and Karpathos (KK). While this may be true for the majority of the Kikladic islands, our results demonstrate that Folegandros is phytogeographically closer to KK, as we recorded eight endemic taxa (*Anthyllis splendens* Willd., *Campanula laciniata* L., *Dianthus fruticosus* L. subsp. *amorginus* Runemark, *Eryngium amorginum* Rech.f., *Erysimum candicum* Snogerup subsp. *candicum*, *Galium amorginum* Halácsy, *Pimpinella pretenderis* (Heldr.) Orph. ex Halácsy and *Sternbergia greuteriana* Kamari & Artelari) occurring exclusively in Kik and KK and only two taxa (*Silene cythnia* (Halácsy) Walters and *Campanula calaminthifolia* Lam.) that occur exclusively in Kik and EAe. Therefore we argue that Folegandros seems to be more closely connected to KK, concurring with previous studies in the southern part of the phytogeographical area of the Kiklades (Kougioumoutzis *et al.*, 2012, 2014a). It could be argued that the southern Kiklades as a whole have higher phytogeographical affinities with Kriti and this could possibly be attributed to the close palaeogeographical distance between southern Kiklades and Kriti during the Messinian salinity crisis (Hsü, 1972). Our results concur with Kougioumoutzis *et al.* (2014b), according to whom most of the biregional endemics present in the phytogeographical region of the Kiklades originate from the phytogeographical region of Kriti and Karpathos.

The conservation status of the Greek endemic taxa of Folegandros, as well as their evaluation status within the Natura 2000 Network, is shown in Table 6. Twenty-four out of 40 endemic taxa are under statutory protection.

TABLE 6. Greek endemic taxa in Folegandros, their geographical distribution and their protection and evaluation status according to European and national legislation and lists

Family	Taxon	Pe	StE	WAe	IoI	SPi	NPi	EC	NC	NE	NAe	Kik	KK	E Ae	Protection status	Natura 2000
Apiaceae	<i>Eryngium amorginum</i> Rech.f.											*	*		RDB (R), PD 67/81	—
Apiaceae	<i>Pimpinella pretenderis</i> (Heldr.) Halácsy											*	*		RDB (R), PD 67/81	—
Apiaceae	<i>Seseli gummiferum</i> Sm. subsp. <i>crithmifolium</i> (DC.) P.H.Davis											*	*	*	PD 67/81	—
Asteraceae	<i>Anthemis scopulorum</i> Rech.f.	*		*								*	*	*	—	—
Asteraceae	<i>Centaurea laconica</i> Boiss. subsp. <i>lineariloba</i> (Halácsy) Wagenitz & E.Gamal-Eldin											*			IUCN (R)	—
Asteraceae	<i>Centaurea raphanina</i> Sm. subsp. <i>mixta</i> (DC.) Runemark	*	*	*								*			—	B
Asteraceae	<i>Crepis hellenica</i> Kamari subsp. <i>hellenica</i>	*	*	*		*		*		?		*		*	—	B
Asteraceae	<i>Filago cretensis</i> Gand. subsp. <i>cretensis</i>	*	*	*								*	*	*	—	B
Asteraceae	<i>Scorzonera cretica</i> Willd.	*										*	*	*	—	—
Boraginaceae	<i>Anchusella variegata</i> (L.) Bigazzi, Nardi & Selvi	*	*	*	*			*	?		*	*	*	*	—	—
Brassicaceae	<i>Erysimum candicum</i> Snogerup subsp. <i>candicum</i>		*	*											PD 67/81	—
Brassicaceae	<i>Erysimum senoneri</i> (Heldr. & Sartori) Wettst. subsp. <i>senoneri</i>			*								*			—	B
Brassicaceae	<i>Fibigia lunarioides</i> (Willd.) Sm.											*	*	*	PD 67/81	B
Campanulaceae	<i>Campanula anchusiflora</i> Sm.	*										*			—	—
Campanulaceae	<i>Campanula calaminthifolia</i> Lam.											*		*	—	—
Campanulaceae	<i>Campanula heterophylla</i> L.											*			PD 67/81	—

TABLE 6. (Cont'd)

Campanulaceae	<i>Campanula laciniata</i> L.									*	*		RDB (R), PD 67/81	B
Caryophyllaceae	<i>Arenaria aegaea</i> Rech.f.	*								*	*	*	—	B
Caryophyllaceae	<i>Dianthus fruticosus</i> L. subsp. <i>amorginus</i> Runemark									*	*		PD 67/81	—
Caryophyllaceae	<i>Silene adelphiae</i> Runemark									*			—	—
Caryophyllaceae	<i>Silene cythnia</i> (Halácsy) Walters									*		*	PD 67/81	—
Chenopodiaceae	<i>Atriplex recurva</i> d'Urv.	*	*			*	*			*	*		—	—
Fabaceae	<i>Anthyllis splendens</i> Willd.									*	*		RDB (R), PD 67/81	B
Fabaceae	<i>Vicia cretica</i> Boiss. & Heldr. subsp. <i>aegaea</i> (Halácsy) P.W.Ball	*								*			PD 67/81	—
Plumbaginaceae	<i>Limonium doerfleri</i> (Halácsy) Rech.f.									*	*	*	—	—
Plumbaginaceae	<i>Limonium ocymifolium</i> (Poir.) Kuntze	*	*							*	*	*	—	B
Primulaceae	<i>Cyclamen graecum</i> Link subsp. <i>graecum</i>	*	*	*	*	*	*	*	*	*			CITES	—
Ranunculaceae	<i>Nigella degenii</i> Vierh. subsp. <i>degenii</i>									*			—	—
Ranunculaceae	<i>Nigella doerfleri</i> Vierh.	*								*	*		—	B
Rubiaceae	<i>Galium amorginum</i> Halácsy									*	*		PD 67/81	—
Scrophulariaceae	<i>Scrophularia heterophylla</i> Willd. var. <i>heterophylla</i>	*	*	*	*	*	*	*	*	*	*	*	—	—
Amaryllidaceae	<i>Sternbergia greuteriana</i> Kamari & Artelari									*	*		—	—
Asparagaceae	<i>Bellevalia hyacinthoides</i> (Bertol.) K.M.Perss. & Wendelbo	*	*	*	*	*	*	?		*			—	—

TABLE 6. (Cont'd)

Family	Taxon	Pe	StE	WAe	IoI	SPi	NPi	EC	NC	NE	NAe	Kik	KK	EAe	Protection status	Natura 2000
Asparagaceae	<i>Muscari cycladicum</i> P.H.Davis & D.C.Stuart subsp. <i>cycladicum</i>											*			—	—
Iridaceae	<i>Crocus cartwrightianus</i> Herb.	*	*		?							*	*		—	—
Iridaceae	<i>Crocus laevigatus</i> Bory & Chaub.	*	*	*						?		*	*	*	—	—
Iridaceae	<i>Crocus tournefortii</i> J.Gay	*										*	*	*	—	B
Liliaceae	<i>Fritillaria ehrhartii</i> Boiss. & Orph.			*								*			—	—
Orchidaceae	<i>Ophrys spruneri</i> Nyman subsp. <i>spruneri</i>	*	*	*	*	*	*					*	*	*	CITES	—
Poaceae	<i>Phleum exaratum</i> Griseb. subsp. <i>aegaeum</i> (Vierh.) Doğan	*										*	*	*	—	—

Abbreviations:

Pe, Peloponnisos; StE, Sterea Hellas; WAe, West Aegean Islands; IoI, Ionian Islands; SPi, South Pindhos; NPi, North Pindhos; EC, East Central; NC, North Central; NE, North East; NAe, North Aegean Islands; Kik, Kiklades; KK, Kriti and Karpathos; EAe, East Aegean Islands.

CITES: Convention on International Trade in Endangered Species of Wild Fauna and Flora.

IUCN: *Red List of Threatened Plants* (IUCN, 2010), with the following classification system: R: the species population is rare.

Natura 2000 (Dafis *et al.*, 1996): The database created after the Directive 43/1992, where the plants are evaluated as: B: Greek endemics.

PD: Greek Presidential Decree 67/1981 (1981), on the protection of the native flora and wild fauna of Greece.

RDB: *Red Data Book of Rare and Threatened Plants of Greece*.

Phytogeographical relationships within the Kiklades

The phytogeographical region of the Kiklades consists of 22 large islands; detailed and available floristic data exist, however, only for the following islands: Anafi, Andros, Antiparos and Paros, Giaros, Kimolos, Milos, Naxos, Santorini and Serifos. Therefore, we focus on these 10 islands in order to examine the floristic affinities of Folegandros Island within the phytogeographical region of the Kiklades.

All the islands in the comparison are in the same bioclimatic zone and phytogeographical region (Kik) as the study area, except for Naxos, which has a more humid climate, although being part of the same phytogeographical region (Kik).

In Table 7 Sørensen's index values for each island pair show that Folegandros has the strongest phytogeographical affinity with Anafi.

DISCUSSION

The high percentages of therophytes (57.48%) and of leguminous taxa (13.43%) indicate disturbance in Mediterranean ecosystems (Naveh, 1974; Arianoutsou & Margaris, 1981; Barbero *et al.*, 1990; Panitsa *et al.*, 1994, 2003; Panitsa & Tzanoudakis, 1998). Intense stock farming and other agricultural activities have not yet ceased in Folegandros and have clearly altered the island's floristic character, as the relatively high local amount of Asian elements (8.55%) suggests.

According to Arianoutsou *et al.* (2010), the total number of alien taxa accounts for c.5% of the native flora of Greece, which is significantly higher than that of Folegandros (1.27%). Nevertheless, in Folegandros where abandoned grazing grounds and farm lands occupy large areas, *Aeonium arboreum* has heavily contaminated and altered these habitats which would otherwise be colonised by native pioneer herbs and shrubs. This phenomenon is also observed in other Aegean islands (Arianoutsou *et al.*, 2010; Kougioumoutzis *et al.*, 2012, 2014a).

TABLE 7. Sørensen's index values for each large island of the phytogeographical region of the Kiklades compared to Folegandros Island

Pair with Folegandros	Sørensen's index
Anafi	63.4
Kimolos	55.4
Santorini	54.7
Antiparos	54.4
Paros	53.1
Serifos	53.0
Milos	52.4
Naxos	47.3
Andros	43.9
Giaros	43.1

The high percentages of chamaephytes and hemicryptophytes depend on the frequency of limestone cliffs which very often harbour endemic taxa (Kypriotakis, 1998; Kypriotakis & Tzanoudakis, 2001; Tzanoudakis *et al.*, 2006). In Folegandros, nearly half (45%) of the endemic flora are chamaephytes or hemicryptophytes, which are scattered in the numerous steep calcareous cliffs present on the island.

The number of species per surface unit is an important parameter of Aegean vascular plant diversity, in relation to the conservation of the diversity of the Aegean area (Panitsa & Tzanoudakis, 2010). In this context Folegandros seems to be a biodiversity hotspot at least for the phytogeographical region of the Kiklades, in spite of the intense human presence and the relatively low number of plant taxa present on the island, as it hosts more than five times (14.7 species/km²) the taxa per surface unit than Andros or Naxos (2.7 and 2.4 species/km², respectively), 37 times the taxa compared to the whole East Aegean area (0.4 species/km², Panitsa & Tzanoudakis, 2010) and 27 times the taxa compared to the Kiklades (c.0.54 species/km², Phitos *et al.*, 1995). Moreover, Folegandros hosts 10 times (1.25 endemic species/km²) the endemic taxa per surface unit compared to Andros and Naxos (0.11 and 0.12 endemic species/km², respectively) and 21 times the endemic taxa compared to the Kiklades (0.06 endemic species/km²). Our results concur with Kougioumoutzis & Tiniakou (2014), who characterised Folegandros as an endemic plant diversity hotspot in the phytogeographical region of the Kiklades.

The existence of biregional endemics is a good indication of phytogeographical connections between regions (Georghiou & Delipetrou, 2010). Eight endemic taxa found in the study area – *Anthyllis splendens*, *Campanula laciniata*, *Dianthus fruticosus* subsp. *amorginus*, *Eryngium amorginum*, *Erysimum candicum* subsp. *candicum*, *Galium amorginum*, *Pimpinella pretenderis* and *Sternbergia greuteriana* – provide useful information regarding the biogeographical position of Folegandros, as they are found exclusively in the phytogeographical regions of Kiklades and Kriti-Karpathos. Folegandros seems to have very close phytogeographical affinities with Amorgos, Anafi and Astypalaea. More specifically, *Erysimum candicum* subsp. *candicum* and *Sternbergia greuteriana* can only be found outside the phytogeographical region of Kriti-Karpathos, in Anafi and Folegandros. The rest of the aforementioned taxa have a wider southern Kikladic distribution; most of them, however, are also found in (or were described from) Amorgos. In particular, *Anthyllis splendens* is found outside Kriti only in Amorgos, Anafi, Folegandros and Kythnos, *Campanula laciniata* is reported from Kriti, Karpathos, Amorgos, Anafi, Astypalaea, Folegandros, Kythnos, Sikinos and Sifnos, *Dianthus fruticosus* subsp. *amorginus* is distributed in Kriti and Amorgos, Astypalaea, Folegandros, Kimolos, Milos and Sikinos, *Eryngium amorginum* is found in Kriti, Amorgos, Astypalaea and Folegandros, *Galium amorginum* in Kriti, Amorgos, Folegandros and Naxos, while *Pimpinella pretenderis* is distributed in Kriti, Astypalaea, Folegandros, Ios, Santorini and Sikinos. The evidence presented here suggests a close phytogeographical relationship between Folegandros and Kriti, as well as between southern Kiklades and Kriti since, according to Strid & Tan (1997),

the phytogeographical region of Kriti and Karpathos has strong connections to that of the Kiklades, especially as far as the dry southeastern islands are concerned.

The flora of Folegandros is more similar to that of Anafi, and then to Kimolos and Santorini (Kik), than to that of Andros and Giaros. The stronger floristic affinities between Folegandros and Anafi can be attributed to the fact that in both these islands there are numerous calcareous cliffs hosting several rare and endemic taxa, as well as to their quite similar climatic conditions. On the other hand, the rather low floristic similarity between Folegandros and Andros can be attributed to the more humid climate of the latter; after all, the phytogeographical region of the Kiklades is climatically compartmentalised (Theocharatos, 1978). Our results are in accordance with the phytogeographical compartmentalisation of the Kiklades (Kougioumoutzis *et al.*, 2014b), as well as with the fact that Folegandros acts as an endemic species pool in the phytogeographical region of the Kiklades (Kougioumoutzis *et al.*, 2014b).

ACKNOWLEDGEMENTS

The authors are much indebted to Dr Paraskevas Vasilakopoulos for his invaluable assistance in the field. Cordial thanks are also due to Dr Ioannis Kokkoris for his help in constructing the geographical map of Folegandros Island and to Dr Leonardos Tiniakos for his help and comments regarding the geology of Folegandros Island. The authors would like to thank Dr Thomas Raus and an anonymous reviewer for their valuable comments and suggestions on the manuscript.

REFERENCES

- AL-EISAWI, D. & JURY, S. L. (1988). A taxonomic revision of the genus *Tordylium* L. (Apiaceae). *Bot. J. Linn. Soc.* 97: 357–403.
- APG III [THE ANGIOSPERM PHYLOGENY GROUP] (2009). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* 161(2): 105–121.
- ARIANOUSOU, M. & MARGARIS, N. S. (1981). Producers and the fire cycle in a phryganic ecosystem. In: MARGARIS, N. S. & MOONEY, H. A. (eds) *Components of Productivity of Mediterranean Climate Regions: Basic and Applied Aspects*, pp. 181–190. Hague: Dr W. Junk Publications.
- ARIANOUSOU, M., BAZOS, I., DELIPEIROU, P. & KOKKORIS, Y. (2010). The alien flora of Greece: taxonomy, life traits and habitat preferences. *Biol. Invasions* 12(10): 3525–3549.
- BARBERO, M., BONIN, G., LOISEL, R. & QUÉZEL, P. (1990). Changes and disturbances of forest ecosystems caused by human activities in the western part of the Mediterranean basin. *Vegetatio* 87: 151–173.
- BERGMEIER, E. & STRID, A. (2014). Regional diversity, population trends and threat assessment of the weeds of traditional agriculture in Greece. *Bot. J. Linn. Soc.* 175(4): 607–623.
- BIEL, B. & TAN, K. (2008). Reports 16–60. In: VLADIMIROV, V., DANE, F. & KIT, T. (eds) *New floristic records in the Balkans* 7, pp. 134–139. *Phytol. Balcan.* 14.

- BLONDEL, J., ARONSON, J., BODIOU, J. Y. & BOEUF, G. (2010). *The Mediterranean Region: Biological Diversity in Space and Time*. Oxford: Oxford University Press.
- BÖHLING, N. B. (1994). *Studien zur landschaftökologischen Raumgliederung auf der mediterranen Insel Naxos (Griechenland); unter besonderer Berücksichtigung von Zeigerpflanzen*. *Dissertationes Botanicae* 230. Berlin, Stuttgart: J. Cramer Verlag.
- BROWICZ, K. (1997). Woody flora of Milos and Kimolos (Kiklades, Greece). *Arbor. Kornickie* 42: 45–63.
- BRULLO, S. & DE MARCO, G. (2000). Taxonomical revision of the genus *Dittrichia* (Asteraceae). *Portugaliae Acta Biol.* 19: 341–354.
- DAFIS, S., PAPASTERGIADOU, E., GEORGHIOU, K., BABALONAS, D., GEORGIADIS, TH., PAPAGEORGIOU, M., LAZARIDOU, TH. & TSIAOUSSI, B. (eds) (1996). *Directive 92/42/EEC. The Greek Habitat Project Natura 2000. An overview*. Thessaloniki (in Greek).
- DAVIS, P. H. (1965–1985). *Flora of Turkey and the East Aegean Islands*. Vols 1–9. Edinburgh: Edinburgh University Press.
- DIMOPOULOS, P., RAUS, TH., BERGMEIER, E., CONSTANTINIDIS, TH., IATROU, G., KOKKINI, S., STRID, A. & TZANOUDAKIS, D. (2013). *Vascular Plants of Greece: An Annotated Checklist*. Berlin (Germany): Botanic Garden and Botanical Museum Berlin-Dahlem. Athens (Greece): Hellenic Botanical Society.
- GALLEGO, M. J., TALAVERA, S. & SILVESTRE, S. (1980). Revision del genero *Reichardia* Roth (Compositae). *Lagascalia* 9(2): 159–217.
- GEORGHIOU, K. & DELIPETROU, P. (2010). Patterns and traits of the endemic plants of Greece. *Bot. J. Linn. Soc.* 162: 130–422.
- GEORGIU, O. (1990). *Biosistimatiki meleti tis omadas Anthemis tomentosa (Asteraceae) stin Ellada* (in Greek). PhD thesis, University of Patras.
- GOUVAS, M. & SAKELLARIOU, N. (2011). *Climate and Forest Vegetation of Greece*. Athens: Institute of Environmental Research and Sustainable Development, National Observatory of Athens.
- GREUTER, W. & RAAB-STRAUBE, E. VON (2008). *Med-Checklist*. Vol. 2. Palermo: Genève & Berlin.
- GREUTER, W., BURDET, M. & LONG, G. (1984–1989). *Med-Checklist*. Vols 1, 3, 4. Genève & Berlin.
- HANSEN, A. (1971). Flora der Inselgruppe Santorin. *Candollea* 26: 109–163.
- HSÜ, K. J. (1972). Origin of saline giants: a critical review after the discovery of the Mediterranean evaporates. *Earth-Science* 8: 371–396.
- IUCN (2010). *IUCN Red List of Threatened Species*. Version 2010.4.
- KAMARI, G. (1976). *Kyttarotaxinomiki meleti tis omadas Crepis neglecta en Elladi* (in Greek). PhD thesis, University of Patras.
- KOUGIOUMOUTZIS, K. & TINIAKOU, A. (2014). Ecological factors and plant species diversity in the South Aegean Volcanic Arc and other central Aegean Islands. *Pl. Ecol. Divers.* doi: 10.1080/17550874.2013.866989.
- KOUGIOUMOUTZIS, K., TINIAKOU, A., GEORGIU, O. & GEORGIADIS, TH. (2012). Contribution to the flora of the South Aegean Volcanic Arc: Anafi Island (Kiklades, Greece). *Willdenowia* 42: 127–141.
- KOUGIOUMOUTZIS, K., TINIAKOU, A., GEORGIU, O. & GEORGIADIS, TH. (2014a). Contribution to the flora of the South Aegean Volcanic Arc: Kimolos Island (Kiklades, Greece). *Edinburgh J. Bot.* 71(2): 135–160.
- KOUGIOUMOUTZIS, K., SIMAIAKIS, S. M. & TINIAKOU, A. (2014b). Network biogeographical analysis of the central Aegean archipelago. *J. Biogeogr.* 41(10): 1848–1858.
- KYPRIOTAKIS, Z. (1998). *Contribution to the chasmophytic flora of Crete*. PhD thesis, University of Patras.

- KYPRIOTAKIS, Z. & TZANOUDAKIS, D. (2001). Contribution to the study of the Greek insular flora: the chasmophytic flora of Crete. *Boccone* 13: 495–503.
- LIVANIOU-TINIAKOU, A., CHRISTODOULAKIS, D., GEORGIIOU, O. & ARTELARI, R. (2003). Floristic dynamics in correlation with the type of substrate and human activities: the example of Serifos (Kyklades Islands, Greece). *Fresenius Environm. Bull.* 12: 1520–1529.
- MÉDAIL, F. & QUÉZEL, P. (1997). Hot-spots analysis for conservation of plant biodiversity in the Mediterranean basin. *Vegetatio* 98: 1–12.
- NAVEH, Z. (1974). Effects of fire in the Mediterranean region. In: KOZŁOWSKI, T. T. & AHLGREN, C. E. (eds) *Fire and Ecosystems*, pp. 401–434. New York: Academic Press.
- PANITSA, M. & TZANOUDAKIS, D. (1998). Contribution to the study of the Greek flora: flora and vegetation of the islands Agathonisi and Pharmakonisi (East Aegean area, Greece). *Willdenowia* 28: 95–116.
- PANITSA, M. & TZANOUDAKIS, D. (2010). Floristic diversity on small islands and islets: Leros islets' group (East Aegean area, Greece). *Phytol. Balcan.* 16(2): 271–284.
- PANITSA, M., DIMOPOULOS, P., IATROU, G. & TZANOUDAKIS, D. (1994). Contribution to the study of the Greek flora: vegetation of the Enousses (Oinousses) islands (E Aegean area). *Flora* 189: 367–374.
- PANITSA, M., SNOGERUP, B., SNOGERUP, S. & TZANOUDAKIS, D. (2003). Floristic investigation of Lemnos island (NE Aegean area, Greece). *Willdenowia* 33: 79–105.
- PAPANIKOLAOU, D. (1984). The three metamorphic belts of the Hellenides: a review and a kinematic interpretation. In: DIXON, J. E. & ROBERTSON, A. H. F. (eds) *The geological evolution of the Eastern Mediterranean. Geological Society of London, Special Publications* 17: 551–561.
- PAPANIKOLAOU, D. (1987). Tectonic evolution of the Cycladic blueschist belt (Aegean Sea, Greece). In: HELGESON, H. C. (ed.) *Chemical Transport in Metasomatic Processes*, pp. 429–450. Dordrecht: Reidel Publishing Company.
- PHITOS, D., STRID, A., SNOGERUP, S. & GREUTER, W. (1995). *The Red Data Book of Rare and Threatened Plants of Greece*. Athens: WWF.
- PHOTIADES, A. & KEAY, S. (2003). Geological and geochronological data for Sikinos and Folegandros metamorphic units (Cyclades, Greece): their tectono-stratigraphic significance. *Bull. Geol. Soc. Greece* 35: 35–45.
- PIGNATTI, S. (1982). *Flora d' Italia*. Bologna.
- PODLECH, D. (2008). The genus *Astragalus* L. (Fabaceae) in Europe with exclusion of the former Soviet Union. *Feddes Repert.* 119: 310–387.
- RAUS, T. (1986). Flora von Paros und Antiparos (Kykladen, Griechenland). *Ann. Naturhist. Mus. Wien* 98: 237–278.
- RAUS, T. (1988). Vascular plant colonization and vegetation development on sea-born volcanic islands in the Aegean (Greece). *Vegetatio* 77: 139–147.
- RAUS, T. (2012). Gefäßpflanzen von Milos (Kykladen, Griechenland) – eine floristische Handreichung. *Verh. Zool.-Bot. Ges. Österreich* 148/149: 197–235.
- RECHINGER, K. H. (1944 [1943]). Flora Aegaea. *Akad. Wiss. Wien, Math.-Naturwiss. Kl., Denkschr.* 105(1): 1–924.
- RUNEMARK, H. (1996). Reports 590–678. In: KAMARI, G., FELBER, F. & GARBARI, F. (eds) *Mediterranean chromosome reports* 6, pp. 223–243. *Fl. Medit.* 6.
- RUNEMARK, H. (2000). Reports 1110–1188. In: KAMARI, G., FELBER, F. & GARBARI, F. (eds) *Mediterranean chromosome reports* 10, pp. 386–402. *Fl. Medit.* 10.
- RUNEMARK, H. (2006). Reports 1473–1571. In: KAMARI, G., FELBER, F. & GARBARI, F. (eds) *Mediterranean chromosome reports* 16, pp. 408–425. *Fl. Medit.* 16.
- SELVI, F. & BIGAZZI, M. (2003). Revision of genus *Anchusa* (Boraginaceae–Boraginae) in Greece. *Bot. J. Linn. Soc.* 142: 431–454.

- SNOGERUP, S. (1994). Reports 267–284. In: KAMARI, G., FELBER, F. & GARBARI, F. (eds) *Mediterranean chromosome reports* 4, pp. 254–258. *Fl. Medit.* 4.
- SNOGERUP, S., SNOGERUP, B., STAMATIADOU, E., VON BOTHMER, R. & GUSTAFSSON, M. (2006). Flora and vegetation of Andros, Kikladhes, Greece. *Ann. Mus. Goulandris* 11: 85–270.
- SØRENSEN, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species content. *Biol. Skr. Kongel. Danske Vidensk. Selsk.* 5(2): 1–34.
- STRID, A. (1996). Phytogeographia Aegaea and the Flora Hellenica Database. *Ann. Naturhist. Hofmus. Wien* 98: 279–289.
- STRID, A. & TAN, K. (1997). *Flora Hellenica*. Vol. 1. Königstein: Koeltz Scientific Books.
- STRID, A. & TAN, K. (2002). *Flora Hellenica*. Vol. 2. Ruggell: A. R. G. Gantner.
- TAN, K. & IATROU, G. (2001). *Endemic Plants of Greece: The Peloponnese*. Copenhagen: Gads Publishers Ltd.
- THANOPOULOS, R. (2007). The genus *Medicago* in Greece: 1. A review of species diversity, geographical distribution and ecological adaptation. *Fl. Medit.* 17: 217–276.
- THEOCHARATOS, G. (1978). *The climate of Cyclades*. PhD thesis, University of Athens, Greece.
- TRIANITIS, K. A. & MYLONAS, M. (2009). Greek islands, biology. In: GILLESPIE, R. & GLAGUE, D. A. (eds) *Encyclopedia of Islands*, pp. 388–392. Berkeley: University of California Press.
- TZANOUDAKIS, D. (1981). Contribution to the study of Cyclades flora: the flora of the island of Yiaros. *Bot. Chron.* 1: 124–136.
- TZANOUDAKIS, D., PANITSA, M., TRIGAS, P. & IATROU, G. (2006). Floristic and phytosociological investigation of the island Antikythera and nearby islets (SW Aegean, Greece). *Willdenowia* 36: 285–301.
- TURRILL, W. B. (1929). *The Plant-Life of the Balkan Peninsula*. Oxford.
- TUTIN, T. G., HEYWOOD, V. H., BURGESS, N. A., MOORE, D. M., VALENTINE, D. H., WALTERS, S. M. & WEBB, D. A. (1964–1980, 1993). *Flora Europaea*. Vols 1–5. Cambridge, UK: Cambridge University Press.
- ZOHARY, M. & HELLER, D. (1984). *The Genus Trifolium*. Jerusalem: Ahva Printing Press.

*Received 25 November 2014; accepted for publication 17 February 2015;
first published online 21 May 2015*

APPENDIX

Notes

- Only taxa new to the investigated area appear in the catalogue below.
- Names of taxa not native to the area are in square brackets.

Abbreviations used

KK: K. Kougioumoutzis observations and/or vouchers
 Obs.: Field observation
 Phot.: Photograph

Collection dates

a: 24/03/2013–27/03/2013
b: 20/04/2013–22/04/2013

c: 18/05/2013–19/05/2013
d: 24/11/2013–25/11/2013

Collection sites

1. 109 m. N36°35'46.5" E24°56'4.5"
2. 84 m. N36°35'41.4" E24°56'44.4"
3. 5 m. N36°36'25.6" E24°57'8.4"
4. 388 m. N36°37'15.3" E24°54'40.6"
5. 108 m. N36°37'49.1" E24°54'52.3"
6. 76 m. N36°38'3.2" E24°54'18.0"
7. 152 m. N36°39'10.2" E24°52'27.8"
8. 2 m. N36°37'49.9" E24°51'20.9"
9. 309 m. N36°37'47.8" E24°55'28.8"
10. 226 m. N36°37'24.5" E24°56'4.6"
11. 178 m. N36°37'34.1" E24°54'6.9"
12. 141 m. N36°37'32.7" E24°53'58.8"
13. 106 m. N36°35'44.8" E24°56'12.5"

14. 82 m. N36°35'39.1" E24°56'46.7"
15. 5 m. N36°36'25.6" E24°57'8.4"
16. 136 m. N36°37'24.1" E24°56'12.0"
17. 15 m. N36°37'1.8" E24°50'47.7"
18. 175 m. N36°37'24.8" E24°55'44.4"
19. 159 m. N36°37'41.2" E24°54'58.5"
20. 92 m. N36°35'49.0" E24°56'19.8"
21. 109 m. N36°35'35.9" E24°56'44.5"
22. 379 m. N36°37'15.7" E24°54'39.9"
23. 149 m. N36°37'30.9" E24°54'2.9"
24. 241 m. N36°37'36.4" E24°55'23.3"
25. 274 m. N36°37'48.4" E24°55'29.3"

Life forms

Chamaephyte (Ch)
Hemicryptophyte (H)
Hydrophyte (A)

Geophyte (G)
Phanerophyte (P)
Therophyte (T)

*Chorological groups***Widely distributed taxa**

European-SW Asian (EA)
Euro-Siberian (ES)
Paleotemperate (Pt)
Circumtemperate (Ct)
Subtropical-Tropical (ST)
Cosmopolitan (Co)

Mediterranean taxa

Eastern Mediterranean (EM)
Mediterranean (Me)
Mediterranean-Atlantic (MA)
Mediterranean-European (ME)
Mediterranean-SW Asian (MS)

Balkan taxa

Balkan (Bk)
Balkan-Italian (BI)
Balkan-Anatolian (BA)

Endemic taxa (Endemic)**Adventive taxa (Adv.)**

Catalogue of plants

FERNS**Pteridaceae**

Adiantum capillus-veneris L. – G, ST; L; 17, b, KK 3204

ANGIOSPERMAE**Acanthaceae**

Acanthus spinosus L. – H, Me; KK Phot.

Amaryllidaceae

Allium neapolitanum Cirillo – G, Me; 9, a, KK 2899

Sternbergia greuteriana Kamari & Artelari – G, Endemic; 25, d, KK 3209

Apiaceae

Daucus guttatus Sm. subsp. *guttatus* – T, Me; 13, b, KK 3282; 16, b, KK 3298

Foeniculum vulgare Mill. – H, Me; L; KK Obs.

Prangos ferulacea (L.) Lindl. – H, EA; L; KK Obs.

Scandix pecten-veneris L. – T, EA; L; 1, a, KK 2980; 10, a, KK 3222

Asparagaceae

Muscari commutatum Guss. – G, BI; 4, a, KK 3210

Muscari cycladicum P.H.Davis & D.C.Stuart subsp. *cycladicum* – G, Endemic; 1, a, KK 2896; 5, a, KK 3115

Ornithogalum narbonense L. – G, Me; L; 13, b, KK 3254

Asteraceae

Anthemis rigida (Sm.) Heldr. – T, EM; 7, a, KK 3368

Calendula arvensis (Vaill.) L. – T, ME; L; 2, a, KK 3058; 4, a, KK 2862

Carduus pycnocephalus L. subsp. *pycnocephalus* – T, Me; 1, a, KK 3001

Crepis foetida L. subsp. *rhoeadifolia* (M.Bieb.) Čelak. – T, EA; L; 1, a, KK 2951

Crepis sancta (L.) Bornm. – T, EA; 1, a, KK 2969

Hedypnois rhagadioloides (L.) F.W.Schmidt subsp. *tubaeformis* (Ten.) Hayek – T, Me; L; 1, a, KK 2946; 13, b, KK 3253; 15, b, KK 3330

Hyoseris scabra L. – T, Me; 9, a, KK 2885; 11, a, KK 2850

Hypochaeris cretensis (L.) Bory & Chaub. – T, Me; 13, b, KK 3287; 16, b, KK 3294

Lactuca acanthifolia (Willd.) Boiss. – H, EM; F; KK Obs.

Notobasis syriaca (L.) Cass. – T, Me; 16, b, KK 3299

Ptilostemon chamaepeuce (L.) Less. – H, EM; L; KK Obs.

Rhagadiolus stellatus (L.) Gaertn. – T, Me; 1, a, KK 2978; 13, b, KK 3250

Scorzonera mollis M.Bieb. – H, EA; L; 1, a, KK 2934; 2, a, KK 3063; 11, a, KK 2828; 12, b, KK 3341; 13, b, KK 3255

Sonchus tenerrimus L. – T, Me; L; 3, a, KK 3078; 7, a, KK 2825; 9, a, KK 2912

Tragopogon dubius Scop. – H, EA; 10, a, KK 3219

Tragopogon porrifolius L. – H, Me; 1, a, KK 2922; 11, a, KK 2844; 12, b, KK 3342

Boraginaceae

Anchusa aegyptiaca (L.) DC. – H, EM; L; 1, a, KK 2924

Anchusa italica Retz. – H, Me; KK Obs.

Buglossoides arvensis (L.) I.M.Johnst. – T, Me; 4, a, KK 2871

Cynoglossum columnae Ten. – T, Me; 1, a, KK 2944

Echium diffusum Sm. – T, Me; L; 4, a, KK 2870

Echium parviflorum Moench – T, Me; L; 16, b, KK 3293
Echium plantagineum L. – T, ME; 12, a, KK 2858
Neatostema apulum (L.) I.M.Johnst. – T, Me; L; 13, b, KK 3277
Onosma graecum Boiss. – T, EM; 1, a, KK 2971

Brassicaceae

Erysimum candicum Snogerup subsp. *candicum* – Ch, Endemic; F; 5, a, KK 3103; 6, a, KK 3112; 7, a, KK 2827; 9, a, KK 2883
Matthiola incana (L.) R.Br. – Ch, ME; 17, b, KK 3338

Campanulaceae

Campanula erimus L. – T, ME; L; 13, b, KK 3263

Caryophyllaceae

Spergularia diandra (Guss.) Boiss. – T, EA; L; 7, a, KK 2853; 9, a, KK 2873
Spergularia media (L.) C.Presl – Ch, Me; 15, b, KK 3321

Chenopodiaceae

Salsola tragus L. – T, Pt/[Co]; KK Obs.

Convolvulaceae

Cuscuta palaestina L. – T, Me; L; 14, b, KK 3317

Crassulaceae

[*Aeonium arboreum* (L.) Webb & Berthel.] – H/C, [NW-Afr.]; KK Obs.

Cucurbitaceae

Bryonia cretica L. – G, EM; L; 9, a, KK 2909

Euphorbiaceae

Euphorbia exigua L. – T, ME; L; 2, a, KK 3048
Euphorbia helioscopia L. – T, Co; 1, a, KK 2964
Euphorbia peplus L. – T, Co; 1, a, KK 3009

Fabaceae

Astragalus echinatus Murray – T, Me; L; 15, b, KK 3307
Astragalus hamosus L. – T, MS; L; 6, a, KK 3116
Astragalus spruneri Boiss. – T, Bk; L; 6, a, KK 3193
Coronilla scorpioides (L.) W.D.J.Koch – T, ME; L; 2, a, KK 3036
Hedysarum spinosissimum L. – G, Me; 1, a, KK 3000; 13, b, KK 3258; 15, b, KK 3319
Hippocrepis unisiliquosa L. – T, Me; 11, a, KK 2838
Lathyrus aphaca L. – T, MS; L; 6, a, KK 3199
Lathyrus cicera L. – T, MS; L; 11, a, KK 2837; 13, b, KK 3333
Lathyrus clymenum L. – T, Me; L; 2, a, KK 3064
Lathyrus setifolius L. – T, Me; L; 12, b, KK 3367
Lotus ornithopodioides L. – T, Me; L; 1, a, KK 3011; 2, a, KK 3060; 6, a, KK 3117; 9, a, KK 2906; 12, b, KK 3370
Medicago disciformis DC. – T, Me; 10, a, KK 3224
Medicago polymorpha L. – T, Pt; 1, a, KK 2953; 6, a, KK 3112
Medicago truncatula Gaertn. – T, MS; L; 2, a, KK 3044; 9, a, KK 2874
Onobrychis aequidentata (Sm.) d'Urv. – T, Me; L; 1, a, KK 2963; 10, a, KK 3218; 15, a, KK 3328
Onobrychis caput-galli Lam. – T, Me; 12, b, KK 3353; 14, b, KK 3301
Ononis ornithopodioides L. – T, Me; 13, b, KK 3267
Ononis viscosa L. subsp. *sieberi* (DC.) Širj. – T, Me; 12, b, KK 3385; 15, b, KK 3308

Trifolium nigrescens Viv. – T, Me; L; 1, a, KK 2935; 9, a, KK 2881

Trifolium spumosum L. – T, Me; L; 7, a, KK 2803

Trifolium tomentosum L. var. *tomentosum* – T, Me; L; 6, a, KK 3105; 7, a, KK 2814; 11, a, KK 2834

Vicia cretica subsp. *aegaea* (Halácsy) P.W.Ball – T, Endemic; 10, a, KK 3225

Vicia hybrida L. – T, Eurymed.; 7, a, KK 2807

Vicia sativa subsp. *cordata* (Hoppe) Batt. – T, ME; 11, a, KK 2863

[*Vicia sativa* L. subsp. *sativa*] – T, Pt; 1, a, KK 2956; 6, a, KK 3123; 11, a, KK 2842; 12, b, KK 3388

Gentianaceae

Centaurium tenuiflorum (Hoffmanns. & Link) Fritsch subsp. *tenuiflorum* – T, ME; 2, a, KK 3205; 12, b, KK 3346; 15, b, KK 3309

Geraniaceae

Erodium gruinum (L.) L'Hér. – T, EM; L; 2, a, KK 3046; 9, a, KK 2908

Geranium molle L. – T, Pt; L; 1, a, KK 2937; 4, a, KK 2864; 16, b, KK 3296

Geranium purpureum Vill. – T, Me; L; KK Obs.

Hypericaceae

Hypericum triquetrifolium Turra – H, EM; L; 1, a, KK 3017

Iridaceae

Crocus cartwrightianus Herb. – G, Endemic; 21, d, KK Phot.; 22, d, KK Phot.

Crocus laevigatus Bory & Chaub. – G, Endemic; 21, d, KK Phot.; 22, d, KK Phot.; 24, d, KK Phot.

Lamiaceae

Lamium moschatum Mill. – T, EM; L; 5, a, KK 3097

Salvia verbenaca L. – H, MA; L; 1, a, KK 2923

Salvia viridis L. – T, Me; L; 1, a, KK 3002; 10, a, KK 3212

Sideritis romana L. subsp. *curvidens* (Stapf) Holmboe – T, EM; L; 11, a, KK 2833

Liliaceae

Gagea graeca (L.) Irmsch. – G, BA; 4, a, KK 2984

Gagea peduncularis (J.Presl & C.Presl) Pascher – G, Me; 4, a, KK 2859

Linaceae

Linum bienne Mill. – H, Me; 2, a, KK 3030

Linum strictum L. subsp. *strictum* – T, Me; 1, a, KK 2995; 12, b, KK 3381; 13, b, KK 3276; 15, b, KK 3315

Malvaceae

Malva parviflora L. – T, MS; L; 7, a, KK 2818

Malva pusilla Sm. – T, ES; 1, a, KK 3013

Myrtaceae

[*Eucalyptus camaldulensis* Dehnh.] – P, [Austr.]; KK Obs.

Oleaceae

Olea europaea L. – P, Me; L; KK Obs.

Orchidaceae

Anacamptis collina (Russell) R.M.Bateman, Pridgeon & M.W.Chase – G, MS; KK Phot.

Anacamptis papilionacea (L.) R.M.Bateman, Pridgeon & M.W.Chase – G, MS; KK Phot.

Ophrys ferrum-equinum Desf. – G, BA; KK Phot.

Ophrys fusca Link subsp. *fusca* – G, Me; KK Phot.

- Ophrys fusca* Link subsp. *iricolor* (Desf.) K.Richt. – G, Me; KK Phot.
Ophrys lutea subsp. *galilaea* (H.Fleischm. & Bornm.) Soó – G, Me; KK Phot.
Ophrys lutea Cav. subsp. *lutea* – G, Me; KK Phot.
Ophrys omegaiifera H.Fleischm. – G, Me; KK Phot.
Ophrys spruneri Nyman subsp. *spruneri* – G, Endemic; KK Phot.
Ophrys tenthredinifera Willd. – G, Me; KK Phot.
Serapias orientalis (Greuter) H.Baumann & Künkele subsp. *orientalis* – G, Me; KK Phot.

Orobanchaceae

- Orobanche minor* Sm. – T, EA; L; KK Phot.
Orobanche pubescens d'Urv. – T, Me; L; KK Phot.
Phelipanche nana (Reut.) Soják – T, Pt; L; KK Phot.

Papaveraceae

- Fumaria capreolata* L. – T, Me; 19, c, KK 3237
Glaucium corniculatum (L.) Rudolph – T, Me; 14, b, KK 3304
Glaucium flavum – H, ME; KK Phot.
Papaver rhoeas L. var. *rhoeas* – T, EM; 1, a, KK 2919; 2, a, KK 3032; 9, a, KK 2890; 11, a, KK 2845; 12, b, KK 3259; 19, c, KK 3234
Papaver rhoeas L. var. *strigosum* Boenn. – T, Pt; 2, a, KK 3019; 13, b, KK 3251

Plantaginaceae

- Plantago afra* L. – T, Me; 2, a, KK 3026; 9, a, KK 2904

Poaceae

- Anisantha diandra* (Roth) Tutin – T, Me; 12, b, KK 3368
Anisantha fasciculata (C.Presl) Nevski – T, Me; 1, a, KK 2993; 13, b, KK 3262
Anisantha sterilis (L.) Nevski – T, MS; L; 13, b, KK 3249
Avena sterilis L. – T, MS; L; 9, a, KK 2877
Briza maxima L. – T, ST; 1, a, KK 2985; 12, b, KK 3248
Hainardia cylindrica (Willd.) Greuter – T, Me; 12, b, KK 3365
Hordeum murinum L. subsp. *leporinum* (Link) Arcang. – T, Me; 2, a, KK 3054; 6, a, KK 3108; 9, a, KK 2914; 12, b, KK 3367; 15, b, KK 3312
Melica minuta L. – H, Me; L; 12, b, KK 3247
Phleum arenarium L. – T, MS; 7, a, KK 2820; 15, b, KK 3326
Phragmites australis (Cav.) Steud. – G, Co; KK Obs.
Piptatherum miliaceum (L.) Coss. – H, MS; 7, a, KK 2810
Poa timoleontis Boiss. – H, EM; 1, a, KK 2926
Polypogon maritimus Willd. subsp. *maritimus* – T, Me; 1, a, KK 2984
Polypogon monspeliensis (L.) Desf. – T, ST; 2, a, KK 3027; 15, b, KK 3313
Stipa capensis Thunb. – T, Me; L; 14, b, KK 3302
Stipa pennata L. – H, ES; 2, a, KK 3047; 13, b, KK 3284

Posidoniaceae

- Posidonia oceanica* (L.) Delile – A, Me; KK Obs.

Primulaceae

- Samolus valerandi* L. – H, Co; 17, b, KK 3200

Rubiaceae

- Galium murale* (L.) All. – T, Me; L; 12, b, KK 3378; 13, b, KK 3260
Galium setaceum Lam. – T, MS; L; 16, b, KK 3290
Galium spurium L. – T, Ct; 1, a, KK 2918; 2, a, KK 3056; 5, a, KK 3095; 7, a, KK 2806; 15, b, KK 3318

Galium verrucosum Huds. – T, Me; 9, a, KK 2895

Sherardia arvensis L. – T, EA; 7, a, KK 2804

Scrophulariaceae

Linaria triphylla (L.) Mill. – T, Me; L; 2, a, KK 3022; 9, a, KK 2891

Scrophularia lucida L. – H (Ch), Me; L; 5, a, KK 3096

Veronica agrestis L. – T, EA; 4, a, KK 2861

Veronica cymbalaria Bodard – T, Me; 9, a, KK 2884

Veronica polita Fr. – T, EA; L; 1, a, KK 2927

Tamaricaceae

Tamarix hampeana Boiss. & Heldr. – NP, EM; 2, a, KK 3023; 7, a, KK 2824

Urticaceae

Parietaria judaica L. – H, EA; 17, b, KK 3336

Valerianaceae

Centranthus calcitrapae (L.) Dufur. – T, Me; L; 2, a, KK 3039; 10, a, KK 3216; 13, b, KK 3283

Valerianella discoidea Loisel. – T, Me; L; 1, a, KK 2929; 2, a, KK 3041

Valerianella vesicaria (L.) Moench – T, MS; 12, b, KK 3390; 13, b, KK 3274